

Facial Weak Order

Aram Dermenjian

Joint work with: Christophe Hohlweg (LACIM) and Vincent Pilaud (CNRS & LIX)

Université du Québec à Montréal

13 July 2016

History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.


History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.
- *Finite Coxeter System* (W, S) such that

$$W := \langle s \in S \mid (s_i s_j)^{m_{i,j}} = e \text{ for } s_i, s_j \in S \rangle$$

where $m_{i,j} \in \mathbb{N}^*$ and $m_{i,j} = 1$ only if $i = j$.

- A *Coxeter diagram* Γ_W for a Coxeter System (W, S) has S as a vertex set and an edge labelled $m_{i,j}$ when $m_{i,j} > 2$.




History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Example

$$W_{B_3} = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^4 = (s_2 s_3)^3 = (s_1 s_3)^2 = e \rangle$$



History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Let (W, S) be a Coxeter system.

- Let $w \in W$ such that $w = s_1 \dots s_n$ for some $s_i \in S$. We say that w has *length* n , $\ell(w) = n$, if n is minimal.
- Let the *(right) weak order* be the order on the Cayley graph where $\overset{w}{\bullet} \xrightarrow{\quad} \overset{ws}{\bullet}$ and $\ell(w) < \ell(ws)$.
- For finite Coxeter systems, there exists a longest element in the weak order, w_0 .


History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Example

Let $\Gamma_{A_2} : s \text{---} t$.

$$sts = w_0 = tst$$



Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers,
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.


Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Parabolic Subgroups

Let $I \subseteq S$.

- $W_I = \langle I \rangle$ is the *standard parabolic subgroup* with long element denoted $w_{0,I}$.
- $W^I := \{w \in W \mid \ell(w) \leq \ell(ws), \text{ for all } s \in I\}$ is the set of minimal length coset representatives for W/W_I .
- Any element $w \in W$ admits a unique factorization $w = w^I \cdot w_I$ with $w^I \in W^I$ and $w_I \in W_I$.
- By convention in this talk xW_I means $x \in W^I$.
- *Coxeter complex* - \mathcal{P}_W - the abstract simplicial complex whose faces are all the standard parabolic cosets of W .



Facial Weak Order

Definition (Krob et.al. [2001], Palacios, Ronco [2006])


The (*right*) *facial weak order* is the order \leq_F on the Coxeter complex \mathcal{P}_W defined by cover relations of two types:

- (1) $xW_I \triangleleft xW_{I \cup \{s\}}$ if $s \notin I$ and $x \in W^{I \cup \{s\}}$,
- (2) $xW_I \triangleleft xw_{0,I}w_{0,I \setminus \{s\}}W_{I \setminus \{s\}}$ if $s \in I$,

where $I \subseteq S$ and $x \in W^I$.

Facial weak order example

- (1) $xW_I \leq xW_{I \cup \{s\}}$ if $s \notin I$ and $x \in W^{I \cup \{s\}}$
 (2) $xW_I \leq xW_{0, I \setminus \{s\}} W_{I \setminus \{s\}}$ if $s \in I$




Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Root System

- Let $(V, \langle \cdot, \cdot \rangle)$ be a Euclidean space.
- Let W be a group generated by a set of reflections S .
 $W \hookrightarrow O(V)$ gives representation as a finite reflection group.
- The reflection associated to $\alpha \in V \setminus \{0\}$ is

$$s_\alpha(v) = v - \frac{2 \langle v, \alpha \rangle}{\|\alpha\|^2} \alpha \quad (v \in V)$$



- A *root system* is $\Phi := \{\alpha \in V \mid s_\alpha \in W, \|\alpha\| = 1\}$
- We have $\Phi = \Phi^+ \sqcup \Phi^-$ decomposable into positive and negative roots.

Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : \bullet \xrightarrow{s} \bullet \xrightarrow{t}$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$



Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : s \text{ --- } t$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$



Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : s \text{ --- } t$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$



Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : s \text{ --- } t$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$



Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : s \text{ --- } t$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$






Weak order and Inversion sets

Given $w, u \in W$ then $w \leq_R u$ if and only if $\mathbf{N}(w) \subseteq \mathbf{N}(u)$.

Example

Let $\Gamma_{A_2} : s \text{---} t$, with Φ given by the roots






Root Inversion Set

Definition (Root Inversion Set)

Let xW_I be a standard parabolic coset. The *root inversion set* is the set

$$\mathbf{R}(xW_I) := x(\Phi^- \cup \Phi_I^+)$$




Note that $N(x) = \mathbf{R}(xW_\emptyset) \cap \Phi^+$.



Root Inversion Set

Example




$$\begin{aligned} \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\ &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\ &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\} \end{aligned}$$



Root Inversion Set

Example




$$\begin{aligned}
 \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\
 &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\
 &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\}
 \end{aligned}$$



Root Inversion Set

Example




$$\begin{aligned}
 \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\
 &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\
 &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\}
 \end{aligned}$$



Root Inversion Set

Example

$$\begin{aligned}
 \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\
 &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\
 &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\}
 \end{aligned}$$






Equivalent definitions

Theorem (D., Hohlweg, Pilaud [2016])




The following conditions are equivalent for two standard parabolic cosets xW_I and yW_J in the Coxeter complex \mathcal{P}_W

- 1 $xW_I \leq_F yW_J$
- 2 $\mathbf{R}(xW_I) \setminus \mathbf{R}(yW_J) \subseteq \Phi^-$ and $\mathbf{R}(yW_J) \setminus \mathbf{R}(xW_I) \subseteq \Phi^+$.
- 3 $x \leq_R y$ and $xw_{0,I} \leq_R yw_{0,J}$.




Equivalence for type A_2 Coxeter System






Equivalence for type A_2 Coxeter System






Equivalence for type A_2 Coxeter System



Equivalence for type A_2 Coxeter System






Equivalence for type A_2 Coxeter System



$$\begin{aligned} \mathbf{R}(xW_I) \setminus \mathbf{R}(yW_J) &\subseteq \Phi^- \\ \mathbf{R}(yW_J) \setminus \mathbf{R}(xW_I) &\subseteq \Phi^+ \end{aligned}$$

$$\begin{aligned} x &\leq_R y \\ xW_{0,I} &\leq_R yW_{0,J} \end{aligned}$$

Equivalence for type A_2 Coxeter System



$$\begin{aligned} \mathbf{R}(xW_I) \setminus \mathbf{R}(yW_J) &\subseteq \Phi^- \\ \mathbf{R}(yW_J) \setminus \mathbf{R}(xW_I) &\subseteq \Phi^+ \end{aligned}$$

$$xW_{0,I} \le_R yW_{0,J}$$

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Facial weak order lattice

Theorem (D., Hohlweg, Pilaud [2016])

The facial weak order (\mathcal{P}_W, \leq_F) is a lattice with the meet and join of two standard parabolic cosets xW_I and yW_J given by:

$$xW_I \wedge yW_J = z_{\wedge} W_{K_{\wedge}},$$

$$xW_I \vee yW_J = z_{\vee} W_{K_{\vee}}.$$

where,



$$z_{\wedge} = x \wedge y \quad \text{and} \quad K_{\wedge} = D_L(z_{\wedge}^{-1}(xw_{0,I} \wedge yw_{0,J})), \text{ and}$$

$$z_{\vee} = xw_{0,I} \vee yw_{0,J} \quad \text{and} \quad K_{\vee} = D_L(z_{\vee}^{-1}(x \vee y))$$

Corollary (D., Hohlweg, Pilaud [2016])

The weak order is a sublattice of the facial weak order lattice.

Example: A_2 and B_2



Example: A_2 and B_2

Example (Meet example)

Recall

$$xW_I \wedge yW_J = z_{\wedge} W_{K_{\wedge}}$$

$$\text{where } z_{\wedge} = x \wedge y$$

$$K_{\wedge} = D_L(z_{\wedge}^{-1}(xw_{o,I} \wedge yw_{o,J}))$$


We compute $ts \wedge stsW_{\{t\}}$.

$$z_{\wedge} = ts \wedge sts = e$$

$$K_{\wedge} = D_L(z_{\wedge}^{-1}(tsw_{o,\emptyset} \wedge stsw_{o,t}))$$

$$= D_L(e(ts \wedge stst))$$

$$= D_L(ts) = \{t\}.$$



Möbius function

Recall that the *Möbius function* of a poset (P, \leq) is the function $\mu : P \times P \rightarrow \mathbb{Z}$ defined inductively by

$$\mu(p, q) := \begin{cases} 1 & \text{if } p = q, \\ - \sum_{p \leq r < q} \mu(p, r) & \text{if } p < q, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition (D., Hohlweg, Pilaud [2016])

The Möbius function of the facial weak order is given by

$$\mu(eW_{\emptyset}, yW_J) = \begin{cases} (-1)^{|J|}, & \text{if } y = e, \\ 0, & \text{otherwise.} \end{cases}$$

Quotients of the facial weak order

Lattice Congruences

Definition

A *lattice congruence* is an equivalence relation \equiv on a lattice (L, \leq) such that for each $x_1 \equiv x_2$ and $y_1 \equiv y_2$ then

- 1 $x_1 \wedge y_1 \equiv x_2 \wedge y_2$, and
- 2 $x_1 \vee y_1 \equiv x_2 \vee y_2$.

Theorem (D., Hohlweg, Pilaud [2016])

Given a lattice congruence \equiv on (W, \leq_R) , the equivalence classes on (\mathcal{P}_W, \leq_F) defined by

$$xW_I \equiv yW_J \iff x \equiv y \text{ and } xw_{o,I} \equiv yw_{o,J}$$

give us a lattice congruence.




Facial Boolean Lattice

Corollary (D., Hohlweg, Pilaud [2016])

Let the (left) *root descent set* of a coset xW_I be the set of roots

$$\mathbf{D}(xW_I) := \mathbf{R}(xW_I) \cap \pm\Delta \subseteq \Phi.$$



Let $xW_I \equiv^{\text{des}} yW_J$ if and only if $\mathbf{D}(xW_I) = \mathbf{D}(yW_J)$.



Facial Cambrian Lattice

Corollary (D., Hohlweg, Pilaud [2016])

Let c be any Coxeter element of W . Let \equiv^c be the c -Cambrian congruence (see Reading [Cambrian Lattice, 2004]). Then let $xW_I \equiv^c yW_J \iff x \equiv^c y$ and $xw_{0,I} \equiv^c yw_{0,J}$.



Thank you!

