

# The facial weak order in hyperplane arrangements

Aram Dermenjian<sup>1,3</sup>

Christophe Hohlweg<sup>1</sup>, Thomas McConville<sup>2</sup> and Vincent Pilaud<sup>3</sup>

<sup>1</sup>Université du Québec à Montréal (UQAM)

<sup>2</sup>Mathematical Sciences Research Institute (MSRI)

<sup>3</sup>École Polytechnique (LIX)

22 April 2019

On this day in 1811 Otto Hesse was born.


# Outline

- Arranging hyperplanes.
- The facial weak order and its 1, 2, 3, 4 (!) definitions.
- Yeah, but is it a lattice?
- Some other properties.

# History and Background - Hyperplanes

- $(V, \langle \cdot, \cdot \rangle)$  -  $n$ -dim real Euclidean vector space.
- A *hyperplane*  $H_i$  is codim 1 subspace of  $V$  with normal  $e_i$ .


## Example



# History and Background - Arrangements


- A *hyperplane arrangement* is  $\mathcal{A} = \{H_1, H_2, \dots, H_k\}$ .
- $\mathcal{A}$  is *central* if  $\{0\} \subseteq \bigcap \mathcal{A}$ .
- Central  $\mathcal{A}$  is *essential* if  $\{0\} = \bigcap \mathcal{A}$ .

## Example



## History and Background - Arrangements

- *Regions*  $\mathcal{R}$  - connected components of  $V$  without  $\mathcal{A}$ .
- *Faces*  $\mathcal{F}_{\mathcal{A}}$  - intersections of closures of some regions.




## History and Background - (Partial) Orders

- *Lattice* - poset where every two elements have a *meet* (greatest lower bound) and *join* (least upper bound).

### Example


The lattice  $(\mathbb{N}, |)$  where  $a \leq b \Leftrightarrow a | b$ .

- meet - greatest common divisor
- join - least common multiple




## History and Background - Poset of regions

- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$



## History and Background - Poset of regions


- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$






## History and Background - Poset of regions

- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$




## History and Background - Poset of regions

- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$



## History and Background - Poset of regions



- *Base region*  $B \in \mathcal{R}$  - some fixed region
- *Separation set for*  $R \in \mathcal{R}$   
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$
- *Poset of Regions*  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  where  
 $R \leq_{\mathcal{A}} R' \Leftrightarrow S(R) \subseteq S(R')$



## History and Background - Poset of regions

- A region  $R$  is *simplicial* if normal vectors for boundary hyperplanes are linearly independent.
- $\mathcal{A}$  is *simplicial* if all  $\mathcal{R}$  simplicial.

### Example






# History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

*If  $\mathcal{A}$  is simplicial then  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice for any  $B \in \mathcal{R}$ . If  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice then  $B$  is simplicial.*

## Example






# History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

*If  $\mathcal{A}$  is simplicial then  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice for any  $B \in \mathcal{R}$ . If  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  is a lattice then  $B$  is simplicial.*

## Example



# Coxeter Arrangements

## Example

A *Coxeter arrangement* is the hyperplane arrangement associated to a Coxeter group.

<b>Coxeter Groups</b>		<b>Hyperplane Arrangements</b>
Reflecting hyperplanes	$\leftrightarrow$	Hyperplane arrangement
Root system	$\leftrightarrow$	Normals to hyperplanes
Inversion sets	$\leftrightarrow$	Separation sets
Weak order	$\leftrightarrow$	Poset of regions

# Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type  $A$ .
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.




# Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type  $A$ .
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.
- Questions: Can we extend this to hyperplane arrangements? Can we find both local and global definitions? When do we actually get a lattice?

# Facial Intervals

Proposition (Björner, Las Vergas, Sturmfels, White, Ziegler '93)

Let  $\mathcal{A}$  be central with base region  $B$ . For every  $F \in \mathcal{F}_{\mathcal{A}}$  there is a unique interval  $[m_F, M_F]$  in  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  such that

$$[m_F, M_F] = \{R \in \mathcal{R} \mid F \subseteq \overline{R}\}$$


# Facial Weak Order



Let  $\mathcal{A}$  be a central hyperplane arrangement and  $B$  a base region in  $\mathcal{R}$ .

## Definition



The *facial weak order* is the order  $\text{FW}(\mathcal{A}, B)$  on  $\mathcal{F}_{\mathcal{A}}$  where for  $F, G \in \mathcal{F}$ :

$$F \leq G \Leftrightarrow m_F \leq_{\mathcal{A}} m_G \text{ and } M_F \leq_{\mathcal{A}} M_G$$



# Facial Weak Order - Example





# Facial Weak Order - Example





# Facial Weak Order - Example



# Facial Weak Order - Example





# Facial Weak Order - Example







# Facial Weak Order - Example



# Facial Weak Order - Example





# Cover Relations

Proposition (D., Hohlweg, McConville, Pilaud, '18+)

For  $F, G \in \mathcal{F}_A$  if

1.  $F \leq G$  in  $\text{FW}(\mathcal{A}, B)$
2.  $|\dim(F) - \dim(G)| = 1$
3.  $F \subseteq G$  or  $G \subseteq F$

then  $F \lessdot G$ .




# Covectors

- *covector* - a vector in  $\{-, 0, +\}^A$  with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$  - set of covectors

## Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$




# Covectors

- *covector* - a vector in  $\{-, 0, +\}^A$  with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$  - set of covectors

## Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$



## Covector operations

For  $X, Y \in \mathcal{L} \subseteq \{-, 0, +\}^{\mathcal{A}}$

■ *Composition*:  $(X \circ Y)(H) = \begin{cases} Y(H) & \text{if } X(H) = 0 \\ X(H) & \text{otherwise} \end{cases}$

■ *Reorientation*:  $(X_{-Y})(H) = \begin{cases} -X(H) & \text{if } Y(H) = 0 \\ X(H) & \text{otherwise} \end{cases}$

★ For  $F \in \mathcal{F}_{\mathcal{A}}$ ,  $[m_F, M_F] = [F \circ B, F \circ -B]$

### Example

Let  $\mathcal{A} = \{H_1, H_2, H_3, H_4, H_5\}$ .

$$X = (-, 0, +, +, 0) \quad Y = (0, 0, -, 0, +)$$

Then

$$X \circ Y = (-, 0, +, +, +) \quad X_{-Y} = (+, 0, +, -, 0)$$





# Covector Definition

## Definition

For  $X, Y \in \mathcal{L}$ :

$$X \leq_{\mathcal{L}} Y \Leftrightarrow Y(H) \leq X(H) \quad \forall H \text{ with } - < 0 < +$$



# Zonotopes

- *Zonotope*  $Z_{\mathcal{A}}$  is the convex polytope:

$$Z_{\mathcal{A}} := \left\{ v \in V \mid v = \sum_{i=1}^k \lambda_i e_i, \text{ such that } |\lambda_i| \leq 1 \text{ for all } i \right\}$$

## Theorem (Edelman '84, McMullen '71)


*There is a bijection between  $\mathcal{F}_{\mathcal{A}}$  and the nonempty faces of  $Z_{\mathcal{A}}$  given by the map*

$$\tau(F) = \left\{ v \in V \mid v = \sum_{F(H_i)=0} \lambda_i e_i + \sum_{F(H_j) \neq 0} \mu_j e_j \right\}$$


*where  $|\lambda_i| \leq 1$  for all  $i$  and  $\mu_j = F(H_j)$*




# Zonotope - Construction example




# Zonotope - Construction example




# Zonotope - Construction example




# Zonotope - Construction example



# Zonotope - Construction example



# Zonotope - Construction example





## Root inversion sets

■ roots  $\Phi_{\mathcal{A}} := \{\pm e_1, \pm e_2, \dots, \pm e_k\}$

■ root inversion set

$$\mathbf{R}(F) := \{e \in \Phi_{\mathcal{A}} \mid \langle x, e \rangle \leq 0 \text{ for some } x \in F\}.$$



# Equivalent definitions

## Theorem (D., Hohlweg, McConville, Pilaud '18+)

For  $F, G \in \mathcal{F}_{\mathcal{A}}$  the following are equivalent:





- $m_F \leq_{\mathcal{A}} m_G$  and  $M_F \leq_{\mathcal{A}} M_G$  in poset of regions  $(\mathcal{R}, B, \leq_{\mathcal{A}})$ .
- There exists a chain of covers in  $\text{FW}(\mathcal{A}, B)$  such that

$$F = F_1 \triangleleft F_2 \triangleleft \cdots \triangleleft F_n = G$$


- $F \leq_{\mathcal{L}} G$  in terms of covectors  $(G(H) \leq F(H) \forall H \in \mathcal{A})$
- $\mathbf{R}(F) \setminus \mathbf{R}(G) \subseteq \Phi_{\mathcal{A}}^-$  and  $\mathbf{R}(G) \setminus \mathbf{R}(F) \subseteq \Phi_{\mathcal{A}}^+$ .



# Equivalence for type $A_2$ Coxeter arrangement





# Equivalence for type $A_2$ Coxeter arrangement




$$m_F \leq_A m_G$$

$$M_F \leq_A M_G$$



$$\mathbf{R}(F) \setminus \mathbf{R}(G) \subseteq \Phi_{\mathcal{A}}^-$$


$$\mathbf{R}(G) \setminus \mathbf{R}(F) \subseteq \Phi_{\mathcal{A}}^+$$



$$F \leq G$$



$$F \subseteq G \text{ or } G \subseteq F$$

$$|\dim F - \dim G| = 1$$





$$G(H) \leq F(H) \quad \forall H \in \mathcal{A}$$






# Equivalence for type $A_2$ Coxeter arrangement







# Equivalence for type $A_2$ Coxeter arrangement




# Equivalence for type $A_2$ Coxeter arrangement







# Equivalence for type $A_2$ Coxeter arrangement



# Equivalence for type $A_2$ Coxeter arrangement



# Equivalence for type $A_2$ Coxeter arrangement





## Facial weak order lattice

Theorem (D., Hohlweg, McConville, Pilaud '18+)

*The facial weak order  $\text{FW}(\mathcal{A}, B)$  is a lattice when  $\mathcal{A}$  is simplicial.*

Corollary (D., Hohlweg, McConville, Pilaud '18+)

*The lattice of regions is a sublattice of the facial weak order lattice when  $\mathcal{A}$  is simplicial.*

# Lattice proof - Joins

Proof uses two key components :


Lemma (Björner, Edelman, Ziegler '90)

*1: If  $L$  is a finite, bounded poset such that  $x \vee y$  exists whenever  $x$  and  $y$  both cover some  $z \in L$ , then  $L$  is a lattice.*


2: Cover relation:  $Z \triangleleft X$  iff  $Z \leq X$ ,  $|\dim X - \dim Z| = 1$  and  $X \subseteq Z$  or  $Z \subseteq X$ . Then  $Z \triangleleft X$  and  $Z \triangleleft Y$  gives three cases:

1.  $X \cup Y \subseteq Z$  and  $\dim X = \dim Y = \dim Z - 1$ ,
2.  $Z \subseteq X \cap Y$  and  $\dim X = \dim Y = \dim Z + 1$ , and
3.  $X \subseteq Z \subseteq Y$  and  $\dim X = \dim Z - 1 = \dim Y - 2$ .


$X \cup Y \subseteq Z$  and  $\dim X = \dim Y = \dim Z - 1$




$X \cup Y \subseteq Z$  and  $\dim X = \dim Y = \dim Z - 1$




$Z \subseteq X \cap Y$  and  $\dim X = \dim Y = \dim Z + 1$




$Z \subseteq X \cap Y$  and  $\dim X = \dim Y = \dim Z + 1$



$X \subseteq Z \subseteq Y$  and  $\dim X = \dim Z - 1 = \dim Y - 2$




$X \subseteq Z \subseteq Y$  and  $\dim X = \dim Z - 1 = \dim Y - 2$







$X \subseteq Z \subseteq Y$  and  $\dim X = \dim Z - 1 = \dim Y - 2$



$X \subseteq Z \subseteq Y$  and  $\dim X = \dim Z - 1 = \dim Y - 2$



## Example: $B_3$ Coxeter arrangement



# Properties of the facial weak order

- The *dual* of a poset  $P$  is the poset  $P^{op}$  where  $x \leq y$  in  $P$  iff  $y \leq x$  in  $P^{op}$ . A poset is *self-dual* if  $P \cong P^{op}$ .
- A lattice is *semi-distributive* if  $x \vee y = x \vee z$  implies  $x \vee y = x \vee (y \wedge z)$  and similarly for the meets.

Theorem (D., Hohlweg, McConville, Pilaud '18+)

*The facial weak order  $\text{FW}(\mathcal{A}, B)$  is self-dual. If furthermore,  $\mathcal{A}$  is simplicial,  $\text{FW}(\mathcal{A}, B)$  is a semi-distributive lattice.*

# Join-irreducible elements

- An element is *join-irreducible* if and only if it covers exactly one element.

Proposition (D., Hohlweg, McConville, Pilaud '18+)

*If  $\mathcal{A}$  is simplicial and  $F$  a face with facial interval  $[m_F, M_F]$ . Then  $F$  is join-irreducible in  $\text{FW}(\mathcal{A}, B)$  if and only if  $M_F$  is join-irreducible in  $(\mathcal{R}, B, \leq_{\mathcal{A}})$  and  $\text{codim}(F) \in \{0, 1\}$*

# Möbius function

Recall that the Möbius function is given by:

$$\mu(x, y) = \begin{cases} 1 & \text{if } x = y \\ -\sum_{x \leq z < y} \mu(x, z) & \text{if } x < y \\ 0 & \text{otherwise} \end{cases}$$


**Proposition (D., Hohlweg, McConville, Pilaud '18+)**

*Let  $X$  and  $Y$  be faces such that  $X \leq Y$  and let  $Z = X \cap Y$ .*

$$\mu(X, Y) = \begin{cases} (-1)^{\text{rk}(X) + \text{rk}(Y)} & \text{if } X \leq Z \leq Y \text{ and } Z = X_{-Z} \cap Y \\ 0 & \text{otherwise} \end{cases}$$

## Further Works

- Can we explicitly state the join/meet of two elements?
- When is the facial weak order congruence uniform?
- Can we generalize this to polytopes?



Thank you!