

Facial Weak Order

Aram Dermenjian

Joint work with: Christophe Hohlweg (LACIM) and Vincent Pilaud (CNRS & LIX)

Université du Québec à Montréal

5 April 2016

History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.


History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.
- *Finite Coxeter System* (W, S) such that

$$W := \langle s \in S \mid (s_i s_j)^{m_{i,j}} = e \text{ for } s_i, s_j \in S \rangle$$

where $m_{i,j} \in \mathbb{N}^*$ and $m_{i,j} = 1$ only if $i = j$.

- A *Coxeter diagram* Γ_W for a Coxeter System (W, S) has S as a vertex set and an edge labelled $m_{i,j}$ when $m_{i,j} > 2$.



History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Example

$$W_{B_3} = \langle s_1, s_2, s_3 \mid s_1^2 = s_2^2 = s_3^2 = (s_1 s_2)^4 = (s_2 s_3)^3 = (s_1 s_3)^2 = e \rangle$$



History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Let (W, S) be a Coxeter system.

- Let $w \in W$ such that $w = s_1 \dots s_n$ for some $s_i \in S$. We say that w has *length* n , $\ell(w) = n$, if n is minimal.
- Let the *(right) weak order* be the order on the Cayley graph where $\overset{w}{\bullet} \xrightarrow{ws} \overset{ws}{\bullet}$ and $\ell(w) < \ell(ws)$.
- For finite Coxeter systems, there exists a longest element in the weak order, w_0 .


History and Background

- The weak order was introduced on Coxeter groups by Björner in 1984, it was shown to be a lattice.

Example

Let $\Gamma_{A_2} : \bullet \xrightarrow{s} \bullet \xrightarrow{t} \bullet$.

$$sts = w_0 = tst$$



Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers,
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and
 - 3 showed that the poset for this order is a lattice.
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.


Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Parabolic Subgroups

Let $I \subseteq S$.

- $W_I = \langle I \rangle$ is the *standard parabolic subgroup* with long element denoted $w_{0,I}$.
- $W^I := \{w \in W \mid \ell(w) \leq \ell(ws), \text{ for all } s \in I\}$ is the set of minimal length coset representatives for W/W_I .
- Any element $w \in W$ admits a unique factorization $w = w^I \cdot w_I$ with $w^I \in W^I$ and $w_I \in W_I$.
- By convention in this talk xW_I means $x \in W^I$.
- *Coxeter complex* - \mathcal{P}_W - the abstract simplicial complex whose faces are all the standard parabolic cosets of W .



Facial Weak Order

Definition (Krob et.al. [2001], Palacios, Ronco [2006])


The (*right*) *facial weak order* is the order \leq_F on the Coxeter complex \mathcal{P}_W defined by cover relations of two types:

- (1) $xW_I \triangleleft xW_{I \cup \{s\}}$ if $s \notin I$ and $x \in W^{I \cup \{s\}}$,
- (2) $xW_I \triangleleft xw_{0,I}w_{0,I \setminus \{s\}}W_{I \setminus \{s\}}$ if $s \in I$,

where $I \subseteq S$ and $x \in W^I$.

Facial weak order example

- (1) $xW_I \leq xW_{I \cup \{s\}}$ if $s \notin I$ and $x \in W^{I \cup \{s\}}$
 (2) $xW_I \leq xw_{o,I}w_{o,I \setminus \{s\}}W_{I \setminus \{s\}}$ if $s \in I$




Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Root System

- Let $(V, \langle \cdot, \cdot \rangle)$ be a Euclidean space.
- Let W be a group generated by a set of reflections S .
 $W \hookrightarrow O(V)$ gives representation as a finite reflection group.
- The reflection associated to $\alpha \in V \setminus \{0\}$ is

$$s_\alpha(v) = v - \frac{2 \langle v, \alpha \rangle}{\|\alpha\|^2} \alpha \quad (v \in V)$$



- A *root system* is $\Phi := \{\alpha \in V \mid s_\alpha \in W, \|\alpha\| = 1\}$
- We have $\Phi = \Phi^+ \sqcup \Phi^-$ decomposable into positive and negative roots.

Inversion Sets


Let (W, S) be a Coxeter system.

Define *(left) inversion sets* as the set $\mathbf{N}(w) := \Phi^+ \cap w(\Phi^-)$.

Example

Let $\Gamma_{A_2} : \underset{s}{\bullet} \text{---} \underset{t}{\bullet}$, with Φ given by the roots

$$\begin{aligned} \mathbf{N}(ts) &= \Phi^+ \cap ts(\Phi^-) \\ &= \Phi^+ \cap \{\alpha_t, \gamma, -\alpha_s\} \\ &= \{\alpha_t, \gamma\} \end{aligned}$$






Weak order and Inversion sets

Given $w, u \in W$ then $w \leq_R u$ if and only if $\mathbf{N}(w) \subseteq \mathbf{N}(u)$.

Example

Let $\Gamma_{A_2} : \bullet \xrightarrow{s} \bullet \xrightarrow{t}$, with Φ given by the roots






Root Inversion Set

Definition (Root Inversion Set)

Let xW_I be a standard parabolic coset. The *root inversion set* is the set

$$\mathbf{R}(xW_I) := x(\Phi^- \cup \Phi_I^+)$$




Note that $N(x) = \mathbf{R}(xW_\emptyset) \cap \Phi^+$.



Root Inversion Set

Example




$$\begin{aligned}
 \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\
 &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\
 &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\}
 \end{aligned}$$



Root Inversion Set

Example

$$\begin{aligned}
 \mathbf{R}(sW_{\{t\}}) &= s(\Phi^- \cup \Phi_{\{t\}}^+) \\
 &= s(\{-\alpha_s, -\alpha_t, -\gamma\} \cup \{\alpha_t\}) \\
 &= \{\alpha_s, -\gamma, -\alpha_t, \gamma\}
 \end{aligned}$$



Equivalent definitions




Theorem (D., Hohlweg, Pilaud [2016])

The following conditions are equivalent for two standard parabolic cosets xW_I and yW_J in the Coxeter complex \mathcal{P}_W




- 1 $xW_I \leq_F yW_J$
- 2 $\mathbf{R}(xW_I) \setminus \mathbf{R}(yW_J) \subseteq \Phi^-$ and $\mathbf{R}(yW_J) \setminus \mathbf{R}(xW_I) \subseteq \Phi^+$.
- 3 $x \leq_R y$ and $xw_{0,I} \leq_R yw_{0,J}$.

Remark Note that showing (1) \Rightarrow (3) and (3) \Rightarrow (2) is easy, but (2) \Rightarrow (1) is more difficult. We used induction on the symmetric difference between the root inversion sets for the proof.

Equivalence for type A_2 Coxeter System



Equivalence for type A_2 Coxeter System



$R(yW_J) \setminus R(xW_I) \subseteq \Phi^+$

$x \leq_R y$
 $xW_{0,I} \leq_R yW_{0,J}$

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order to an order on all faces for type A using inversion tables. They
 - 1 gave a local definition of this order using covers, ✓
 - 2 gave a global definition of this order combinatorially, and ✓
 - 3 showed that the poset for this order is a lattice. ✓
- In 2006, Ronco and Palacios extended this new order to Coxeter groups of all types using cover relations.
- Our motivation was to continue this work for all Coxeter groups.

Facial weak order lattice

Theorem (D., Hohlweg, Pilaud [2016])

The facial weak order (\mathcal{P}_W, \leq_F) is a lattice with the meet and join of two standard parabolic cosets xW_I and yW_J given by:

$$xW_I \wedge yW_J = z_{\wedge} W_{K_{\wedge}},$$

$$xW_I \vee yW_J = z_{\vee} W_{K_{\vee}}.$$

where,



$$z_{\wedge} = x \wedge y \quad \text{and} \quad K_{\wedge} = D_L(z_{\wedge}^{-1}(xw_{0,I} \wedge yw_{0,J})), \text{ and}$$

$$z_{\vee} = xw_{0,I} \vee yw_{0,J} \quad \text{and} \quad K_{\vee} = D_L(z_{\vee}^{-1}(x \vee y))$$

Corollary (D., Hohlweg, Pilaud [2016])

The weak order is a sublattice of the facial weak order lattice.

Example: A_2 and B_2



Example: A_2 and B_2

Example (Meet example)

Recall

$$xW_I \wedge yW_J = z_\wedge W_{K_\wedge}$$

$$\text{where } z_\wedge = x \wedge y$$

$$K_\wedge = D_L(z_\wedge^{-1}(xw_{o,I} \wedge yw_{o,J}))$$


We compute $ts \wedge stsW_{\{t\}}$.

$$z_\wedge = ts \wedge sts = e$$

$$K_\wedge = D_L(z_\wedge^{-1}(tsw_{o,\emptyset} \wedge stsw_{o,t}))$$

$$= D_L(e(ts \wedge stst))$$

$$= D_L(ts) = \{t\}.$$



Proof outline

Recall that $xW_I \leq_F yW_J \Leftrightarrow x \leq_r y$, and $xw_{o,I} \leq_R yw_{o,J}$.

We want to show that $xW_I \wedge yW_J = z_\wedge W_{K_\wedge}$ where $z_\wedge = x \wedge y$ and $K_\wedge = D_L(z_\wedge^{-1}(xw_{o,I} \wedge yw_{o,J}))$

- First we show that this element is in the Coxeter complex $z_\wedge \in W^{K_\wedge}$.
- We then show it's an upper bound: $x \wedge y \leq_R x, y$. Also, $w_{o,K_\wedge} \leq_R z_\wedge^{-1}(xw_{o,I} \wedge yw_{o,J})$ implies $z_\wedge w_{o,K_\wedge} \leq_R xw_{o,I} \wedge yw_{o,J}$.
- Finally we show uniqueness by supposing there exists another element $zW_K \leq_F xW_I, yW_J$. Then we have $z \leq_R x \wedge y = z_\wedge$. Showing $zw_{o,K} \leq_R z_\wedge w_{o,K_\wedge}$ is done by looking at descents and the fact that $z \leq_R z_\wedge$.
- Join is found by an anti-automorphism.

Möbius function

Recall that the *Möbius function* of a poset (P, \leq) is the function $\mu : P \times P \rightarrow \mathbb{Z}$ defined inductively by

$$\mu(p, q) := \begin{cases} 1 & \text{if } p = q, \\ - \sum_{p \leq r < q} \mu(p, r) & \text{if } p < q, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition (D., Hohlweg, Pilaud [2016])

The Möbius function of the facial weak order is given by

$$\mu(eW_{\emptyset}, yW_J) = \begin{cases} (-1)^{|J|}, & \text{if } y = e, \\ 0, & \text{otherwise.} \end{cases}$$

Quotients of the facial weak order

Lattice Congruences

Definition

An *order congruence* is an equivalence relation \equiv on a poset (P, \leq) such that:

- 1 Every equivalence class under \equiv is an interval of (P, \leq) .
- 2 The projection $\pi^\uparrow: P \rightarrow P$ (resp. $\pi^\downarrow: P \rightarrow P$), which maps an element of P to the maximal (resp. minimal) element of its equivalence class, is order preserving.

Theorem (D., Hohlweg, Pilaud [2016])

The equivalence classes, denoted \equiv , given by the following projection give an order congruence on (\mathcal{P}_W, \leq_F) . Let $\Pi^\uparrow(xW_I)$ be the largest standard parabolic coset in $[\pi^\uparrow(x), \pi^\uparrow(xw_{o,I})]$ containing $\pi^\uparrow(x)$.




Facial Boolean Lattice

Corollary (D., Hohlweg, Pilaud [2016])

Let the (left) *root descent set* of a coset xW_I be the set of roots

$$\mathbf{D}(xW_I) := \mathbf{R}(xW_I) \cap \pm\Delta \subseteq \Phi.$$



Let $xW_I \equiv^{\text{des}} yW_J$ if and only if $\mathbf{D}(xW_I) = \mathbf{D}(yW_J)$.



Facial Cambrian Lattice

Corollary (D., Hohlweg, Pilaud [2016])

Let c be any Coxeter element of W . Let \equiv^c be the c -Cambrian congruence (see Reading [Cambrian Lattice, 2004]). Then let $xW_I \equiv^c yW_J \iff x \equiv^c y$ and $xw_{0,I} \equiv^c yw_{0,J}$.



Thank you!

